



## FEVER CONTROL DOSING INFORMATION

### **Infant Tylenol/Acetaminophen Dosing**

**\*\*Only use dosing syringe provided in the bottle for measuring\*\***

6lbs to 11 lbs, use 0.4 ml of Infant Tylenol every 4 hours  
12-17 lbs, use 0.8 ml of Infant Tylenol every 4 hours  
17 ½ lbs-22 lbs, use 1.2 ml (0.8ml + 0.4 ml) every 4 hours  
After 22 lbs, we recommend using Children's Tylenol

### **Children's Tylenol/Acetaminophen Dosing**

22 lbs to under 32 lbs, use 1 tsp (teaspoon) or 5 ml of Children's Tylenol every 4 hours  
32lbs to under 35 lbs, use 1 ¼ tsp every 4 hours  
35 lbs to under 44 lbs, use 1 ½ tsp or 7.5 ml every 4 hours  
44 lbs to under 48 lbs, use 1 ¾ tsp every 4 hours  
48 lbs to 59 lbs, use 2 tsp every 4 hours  
Above 60 lbs, consider Tylenol Meltaway tabs and use instructions on box

### **Infant Motrin/Advil (Ibuprofen) Dosing**

**\*\*Do not use this product below 6 months of age unless advised by physician\*\*\***

**&**

**\*\*Only use dosing syringe provided in the bottle for measuring\*\***

12 lbs to under 17 lbs, use 1.25 ml every 6-8 hours  
17 lbs to under 22 lbs, use 1.875 ml every 6-8 hours  
After 22 lbs, we recommend Children's Motrin/Advil

### **Children's Motrin/Advil (Ibuprofen) Dosing**

22 lbs to under 28 lbs, use 1 tsp every 6-8 hours  
28 lbs to under 33 lbs, use 1 ¼ tsp every 6-8 hours  
33 lbs to under 38 lbs, use 1 ½ tsp every 6-8 hours  
38 lbs to under 44 lbs, use 1 ¾ tsp every 6-8 hours  
44 lbs to 54 lbs, use 2 tsp every 6-8 hours  
55 lbs or higher, consider chewable Motrin tablets and use instructions on box