



**NEW FEVER CONTROL DOSING INFORMATION**

**Infant Tylenol/Acetaminophen Dosing – Concentration (80 mg/0.8 ml)**

**\*\*Only use dosing syringe provided in the bottle for measuring\*\***

6lbs to 11 lbs, use 0.4 ml of Infant Tylenol every 4 hours  
12-17 lbs, use 0.8 ml of Infant Tylenol every 4 hours  
17 ½ lbs-22 lbs, use 1.2 ml (0.8ml + 0.4 ml) every 4 hours  
After 22 lbs, we recommend using the new Infant/Children's Tylenol (160 mg/5ml)

**\*New/Renamed\*Infant/Children's Tylenol/Acetaminophen Dosing (160mg/5ml)**

**\*\*Only use dosing syringe provided in the bottle for measuring\*\***

6 lbs to 11 lbs, use 1.25 ml (1/4 teaspoon) every 4 hours  
12 lbs to 17 lbs, use 2.5 ml (1/2 tsp) every 4 hrs  
18 lbs to 23 lbs, use 3.75 ml (3/4 tsp) every 4 hours  
24 lbs to under 32 lbs, use 5 ml or 1 tsp every 4 hours  
32lbs to 35 lbs, use 6.25 ml or 1 ¼ tsp every 4 hours  
36 lbs to under 44 lbs, use 7.5 ml or 1 ½ tsp every 4 hours  
44 lbs to under 48 lbs, use 8.75 ml or 1 ¾ tsp every 4 hours  
48 lbs to 59 lbs, use 2 tsp every 4 hours  
Above 60 lbs, consider Tylenol Meltaway tabs and use instructions on box

**Infant Motrin/Advil (Ibuprofen) Dosing-Concentration (50 mg/1.25 ml)**

**\*\* Do not use this product below 6 months of age unless advised by physician\*\***

**&**

**\*\*Only use dosing syringe provided in the bottle for measuring\*\***

12 lbs to under 17 lbs, use 1.25 ml every 6-8 hours  
17 lbs to under 22 lbs, use 1.875 ml every 6-8 hours  
After 22 lbs, we recommend Children's Motrin/Advil

**Children's Motrin/Advil (Ibuprofen) Dosing-Concentration (100 mg/5ml)**

**\*\*Do not use this product below 6 months of age unless advised by physician\*\***

**&**

**\*\*Only use dosing syringe provided in the bottle for measuring\*\***

22 lbs to under 28 lbs, use 1 tsp every 6-8 hours  
28 lbs to under 33 lbs, use 1 ¼ tsp every 6-8 hours  
33 lbs to under 38 lbs, use 1 1/2 tsp every 6-8 hours  
38 lbs to under 44 lbs, use 1 ¾ tsp every 6-8 hours  
44 lbs to 54 lbs, use 2 tsp every 6-8 hours  
55 lbs or higher, consider chewable Motrin tablets and use instructions on box